

From: OXYGEN TRANSPORT TO TISSUE - IV
Edited by Haim I. Bicher and Duane F. Bruley
(Plenum Publishing Corporation, 1983)

CHANGES IN TISSUE OXYGENATION AND ACIDITY INDUCED BY LOCALIZED
MICROWAVE HYPERTHERMIA AND HEMATOPORPHYRIN PHOTOTHERAPY, AN UPDATE*

Haim I. Bicher and D. F. Bruley

Western Tumor Medical Group, Valley Cancer Institute -
5522 Sepulveda - Van Nuys, Ca. 91411 - - Louisiana Tech
University - Biomedical Engineering Department - Ruston
Louisiana 71272

INTRODUCTION

Due, in part, to increasing clinical interest, active investigation of the physiological phenomena induced by hyperthermia is in progress (1, 2, 7, 13). The studies of Eddy (7) and Reinhold (13) employing "chamber systems" have both shown changes in the microvascular network as a function of temperature and exposure time. Cater et al. (3) reported on changes in tumor oxygen tension with hyperthermia but did not record changes in tumor temperature. Bicher (1, 2), in a mouse leg tumor system, reported that tumor blood flow increased up to 41°C and then decreased to 44°C. The oxygen tension in the tumor, as measured with a platinum electrode, generally followed the changes in tumor blood flow.

Although blood flow and shifts induced in it by hyperthermia in both tumor and normal tissue is important, several other parameters also have significant roles. Several studies indicate that the pH of interstitial fluid in human and rodent solid tumors is .3 to .5 units lower than the normal tissue pH of about 7.4 (8, 10).

Reduced pH has also been shown to affect the transplantability of tumor cells heated in vitro (12). In a recent paper, Gerweck has shown (9) that there is a variable influence of pH according to temperature and that there is a critical point in the increased lethality of heat below pH 6.7.

*This work was partially reported at the 11th European Conference for Microcirculation, Garmish, 1981 (Bibl.Anat. N-20). Thanks are given to Mr. S. Frinak for technical assistance in part of this work.

In addition to hyperthermia, Hematoporphyrin derivative (HpD) phototherapy is also showing some promise in clinical cancer therapy (5, 6). This type of therapy employs an injectable dye (HpD) which is specifically accumulated in some tumors and/or is specifically cleared from normal tissues (5, 6). When light of specific wavelengths illuminate the dye a photochemical reaction takes place yielding the cytotoxic agent singlet oxygen (7, 15). Although the exact mechanism(s) of tumor inactivation have yet to be determined, the rapid and dramatic coloration changes observed in tumors following treatment suggest that modification of blood flow with possible concomitant effects on pO_2 and pH may play a prominent role (5).

MATERIALS AND METHODS

Animal system. All in situ studies were carried out in 4th, generation transplants of C₃H mammary adenocarcinoma implanted in the hind leg of C₃H SED-BH mice. This is a syngenic implantable tumor that is kept at our facility using solid tissue transplants that are inoculated subcutaneously into recipient mice. Tumors used for experimentation were approximately 8-10mm in diameter. The mice were anesthetized during microelectrode introduction with a combination of Ketamine 40mg/kg I.M. and Thorazine, 50mg/kg I.M.

Oxygen ultramicroelectrodes. The O_2 ultramicroelectrodes used were as described by Cater and colleagues (4). They were made by pulling a glass tube (KG-33, ID 1.5mm, OD 2.0mm, Garner Glass Co., Claremont, California), encasing a 20- μ gold wire (Sigmund Cohn Corp., Mt. Vernon, New York) in a David Kopf Model 700C vertical pipette puller. The exposed gold tip is about 10 μ in diameter, and is coated with a Rhoplex (Rhcm Haas, Philadelphia, Pennsylvania) membrane as previously described (2). This probe is used as an "external reference" O_2 microelectrode.

Electrodes are calibrated as previously described (1) in buffered saline solutions of known pO_2 values. The electrodes are 'conditioned' by placing them in buffered saline and applying 0.8 V potential for 2 hrs. After this treatment they are usually very stable. The current reading at zero oxygen tension is very low (residual current) and the response of the microelectrode to changes of oxygen tension is very rapid.

In these experiments a polarizing voltage of 0.6 V has been used. The relation between current output and oxygen tension is linear, the current per mm Hg being of the order of magnitude of 0.6×10^{-11} A.

pH ultramicroelectrodes. Designs for glass pH microelectrodes employed in this study have been developed, most notably by Hinke (11). The Hinke-type electrode consist of a pH sensitive glass

micropipette inside of pyrex glass pipette with the tip of the pH sensitive micropipette extruded from the pyrex glass pipette. A silver/silver chloride electrode is inserted into the electrode stem which is filled with 0.1 N HCl. This type of microelectrode with an exposed tip has an almost instantaneous response time. This is an advantage over other types of microelectrodes in which a recessed tip may cause a response time of up to several minutes.

Temperature determinations. Tumor and mouse core temperatures were recorded using Copper-Constantan microthermocouples (tip diameter 30-100 microns - MEDTRA Inc.) inserted into the tumoral tissue in close proximity to the microelectrode or in the animal's rectum for core measurements. An Omega Engineering Model 250 Digital Voltmeter amplifier was used as a link between the microthermocouple and the polygraph. Microwaves of a frequency 2450 MHz were produced by a Raytheon Magnetron and delivered through a specially designed 2cm square applicator loaded with low loss dielectric material having a dielectric constant of 6 (14).

HpD phototherapy. Tumor-bearing mice were injected with 20mg/kg HpD and 24 hours later were exposed to the light source. The mice were shielded from the light during the 55 minute exposure except for the tumor-bearing region. The light source was a modified Bessler lantern slide projector which was filtered to yield 150mW/cm² at the tumor surface over the range 600-730nm.

RESULTS

4 different major experiments were performed. In the first pO₂ distribution after 1 hour hyperthermia was studied 1, 4 and 24 hours after treatment. The second experiment was similar but studied the effect of Phototherapy on TpO₂, at similar time intervals. Experiment 3 studied the effect of hyperthermia on tissue pH, at similar time intervals, while experiment 4 determined and compared the effect of Hyperthermia and phototherapy on tissue pH. All results are expressed in Histogram fashion.

Experiment 1: Effect of hyperthermia on TpO₂ - time correlation. Histograms were obtained before and 1, 4 and 24 hours after microwave induced hyperthermia (42-43°C) for one hour. These results are shown in Figure 1. There is a progressive shift towards low pO₂ values, with all the tumor virtually hypoxic after 24 hours. No reoxygenation is noted.

Experiment 2: Effect of Phototherapy on TpO₂ - Time correlation. Histograms were obtained before and 1, 4 and 24 hours after HpD phototherapy (20mg/kg HpD, light exposure 150 mW/cm² for 55 minutes). The pO₂ histogram shifts toward the hypoxic region (0.5 mm Hg) quickly. However, at 4 hours there is an attempt to reoxygenate, as shown by the reappearance of an oxic tail in the




Figure 1. Effect of Hyperthermia on TpO₂.




Figure 2. Effect of HpD Phototherapy on TpO₂.

20-40 mm Hg region. At 24 hours all values are between 0-10 mm Hg. These results are shown in Figure 2.

Experiment 3: Effect of Hyperthermia on T_pH. Time correlation. Histograms obtained before and 1, 4 and 24 hours after microwave reduced Hyperthermia (42-43°C) for one hour. Results shown in Figure 3. Note a remarkable shift of the tissue pH values towards acidity with mean T_pH changing from 6.74 before heat to 6.21 after. T_pH remains in that region at 4 and 24 hours.

Experiment 4: Effect of Hyperthermia and Phototherapy on Tissue pH - a correlation. Histograms obtained before or 1 hour after microwave induced Hyperthermia, Phototherapy or a combination of both. Note that the pH shift towards acidosis (same as shown in figure 3) induced by hyperthermia is not present after Phototherapy. This correlation is shown in Figure 4.

DISCUSSION

Determination of the mode of tumor inactivation by a treatment modality is critical to its development and future use. It has been shown that hyperthermia has many possible effects on cell survival either alone or in combination with radiation (9, 12, 13). It has also been shown to dramatically modify blood flow and oxygenation within tumors (1, 2). The results presented here indicate that a significant reduction in pH is induced by hyperthermia which may result in a significant increase in cell killing within the tumor (9). It is possible that this observed pH shift is due to a combination of hyperthermia stimulated cellular metabolic activity and the simultaneous reduction in tumor blood flow which is observed at the treatment temperature (1).

The effects seen on pH and pO₂ following HpD phototherapy are quite different from those following hyperthermia. It is clear that there is a sharp reduction in pO₂ at all areas within the tumor without any significant shift in pH. Although further studies are now in progress, it is possible to speculate on the meaning of the results presented here. Massive coagulation necrosis within tumors is reported to follow HpD phototherapy (5). It is likely that the cells most affected by this treatment are the vascular endothelial cells of the tumor microvasculature. Their destruction would result in the observed coagulation necrosis with an abrupt reduction in blood flow. This would result in the low levels of tissue oxygenation reported here. Since there is no cellular metabolic stimulation with this modality and there is direct cytotoxicity (15) no dramatic shift in pH would be expected and none was observed.

Further studies are currently in progress to further examine the effects of hyperthermia and HpD phototherapy on tumor




Figure 3. Effect of Hyperthermia on T_{ph}.




Figure 4. Effect of Phototherapy, Hyperthermia or a combination of both on T_{ph}.

MICROPHYSIOLOGY

C₃H Mice Adenocarcinoma Implanted in Leg Muscle

Hyperthermia and Phototherapy

1. O₂ pH inhomogeneities occur in tumor and muscle tissue.
2. Important microcirculation changes occur in both Phototherapy and Hyperthermia
3. One hour 43°C Hyperthermia:
 - A) Collapses Microcirculation.
 - B) O₂ Histogram shifts toward 0-5 mm Hg O₂ Decrease with time.
 - C) No Reoxygenation up to 24 hours.
 - D) pH shifts toward Acidosis.
4. Phototherapy:
 - A) Microcirculation Damage.
 - B) O₂ Histogram shifts toward 0-5 mm Hg quickly.
 - C) Reoxygenation at 4 hours.
 - D) No pH shift.

Figure 5. For explanation see text.

microphysiology. Results obtained so far are summarized in Figure 5.

SUMMARY

Changes in tumor tissue oxygenation and acidity were determined using ultramicroelectrodes, and presented in histogram fashion. The effect of Hyperthermia and Hpd photo-therapy were tested. It was found that both modalities affect tumor microcirculation, causing a marked drop in oxygen availability. Tissue pH is decreased by Hyperthermia, but not by phototherapy. These effects are long lasting, at least for 24 hours after treatment.

REFERENCES

1. Bicher, H. I., Hetzel, F. W., Sandhu, T. S., Frinak, S., Vaupel, P., O'Hara, M. D. and O'Brien, T. Effects of hyperthermia on normal and tumor microenvironment. *Radiology*, In Press 1980.
2. Bicher, H. I. Increase in brain tissue oxygen availability induced by localized microwave hyperthermia. In Silver, Erecinska, and Bicher, *Oxygen transport to tissue*; Vol. III, pp. 347-353, Plenum Press, New York, 1978.
3. Cater, D. B. and Silver, I. A. Quantitative measurements of oxygen tension in normal tissues and in the tumors of patients before and after radiotherapy. *Acta. Radiologica*. 53: 233-256, 1960.

4. Cater, D. B., Silver, I. S. and Wilson, G. M. Apparatus and technique for the quantitative measurement of oxygen tension in living tissues. *Proc. Roy. Soc., Lond. (Series B)* 151: 256-276, 1959-1960.
5. Diamond, I., McDonagh, A. F., Wilson, C. B., Granelli, S. G., Nielsen, S. and Jaenicke, R. Photodynamic therapy of malignant tumors. *Lancet*, 2: 1175-1177, 1972.
6. Dougherty, T. J., Kaufman, J. E., Goldfarb, A., Weishaupf, K. R., Boyle, D., and Mittleman, A. Photoradiation therapy for the treatment of malignant tumors. *Cancer Res.* 38: 2628-2633, 1978.
7. Eddy, H. A. Alterations in tumor microvasculature during hyperthermia. *Radiology*, *In Press*, 1980.
8. Eden, M., and Kahler, H. The pH of rat tumors measured in vivo. *J. Nat'l. Cancer Inst.* 16: 541-556, 1955.
9. Gerweck, L. E. Modification of cell lethality at elevated temperatures: The pH effect. *Radiat. Res.* 70: 224-235 1977.
10. Gullino, P. M., Grantham, F. M. and Smith, S. H. Modifications of the acid-base status of the internal milieu of tumors. *J. Nat'l. Cancer Inst.* 34: 857-860, 1965.
11. Hinke, J. A. Cation-selective microelectrodes for intracellular use. In Eiserman, Glass electrodes for hydrogen and other cations; Marcel Dekker, Inc., New York, 1973.
12. Overgaard, J. Influence of extracellular pH on the viability and morphology of tumor cells exposed to hyperthermia. *J. Nat'l. Cancer Inst.* 56: 1243-1250, 1976.
13. Reinhold, H. S., Blachiewicz, B., and Berg-Blok, A. V. D. Decrease in tumor microcirculation during hyperthermia. in Streffer, *Cancer therapy by hyperthermia and radiation*. pp. 231-232, Urban and Schwarzenburg, Baltimore, 1978.
14. Sandhu, T. S., Kowal, H. S., and Johnson, R. Development of hyperthermia applicators. *Int. J. Radiat. Oncol. Biol. Phys.* 4: 515-519, 1978.
15. Weishaupt, K. R., Gomer, C. J. and Dougherty, T. J. Identification of singlet oxygen as the cytotoxic agent in photo-inactivation of a murine tumor. *Cancer Res.* 36, 2326, 1976.